

OMICS technologies in reproductive medicine: Assessment of quality of oocytes and embryos

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Human Stem Cell Institute. All rights reserved. One of the main factors of success of the procedure art (assisted reproductive technology) is the selection of the most high-quality gametes for further manipulation and obtaining a viable embryo for implantation. The majority of modern techniques based on morphokinetic predictors of quality (i. e. assessment of embryo morphology and rate of division of the blastomeres), which allowed to achieve some success in increasing the percentage of successful pregnancies and reduce the number of multiple pregnancies, but their accuracy is currently insufficient. Thus, the development of objective, reliable, fast and affordable test systems to determine the quality of oocytes and the development potential of the embryo one of the challenges of reproductive medicine. The purpose of this review was to describe the advantages and limitations obecnych technologies, the application of which will allow to deepen our understanding of the physiology of the embryo, as well as set criteria for non-invasive selection of gametes and embryos. In this regard, recently in assisted reproduction are applied the studies of genomic, proteomic, transcript, and metabolomic profiles of oocytes, granulosa and Cumulus cells, embryos, of conditioned media.

<http://dx.doi.org/10.23868/201805003>

Keywords

Blastocyst, Embryo, Genomics, Metabolome, Proteome, Secretome, Transcriptome

References

- [1] Zegers-Hochschild F., Adamson G.D., de Mouzon J., et al. The International Committee for Monitoring Assisted Reproductive Technology and the World Health Organization Revised Glossary on ART Terminology. *Fertility and Sterility* 2009; 92: 1520-4.
- [2] Nimmi R. Review literature on assisted reproductive technology. *International Journal of Current Advanced Research* 2017; 6(2): 2155-6.
- [3] Calhaz-Jorge C., de Geyter C., Kupka M.S., et al. Assisted reproductive technology in Europe, 2012: Results generated from European registers by ESHRE [editorial]. *Human Reproduction* 2016; 31(8): 1638-52.
- [4] Gardner D.K., Meseguer M., Rubio C., et al. Diagnosis of human preimplantation embryo viability. *Hum. Reprod. Update* 2015; 21(6): 727-47.
- [5] Dominguez F., Meseguer M., Aparicio-Ruiz B., et al. New strategy for diagnosing embryo implantation potential by combining proteomics and timelapse technologies. *Fertility and Sterility* 2015; 104(4): 908-14.
- [6] Lai H.H., Chuang T.H., Wong L.K., et al. Identification of mosaic and segmental aneuploidies by next-generation sequencing in preimplantation genetic screening can improve clinical outcomes compared to array-comparative genomic hybridization. *Mol. Cytogenet.* 2017; 10: 14.

- [7] Bromer J., Seli E. Assessment of embryo viability in assisted reproductive technologies: Shortcomings of current approaches and the emerging role of metabolomics. *Current Opinion in Obstetrics and Gynecology* 2008; 20: 234-41.
- [8] Silvestri E., Lombardi A., De Lange P., et al. Studies of complex biological systems with applications to molecular medicine: The need to integrate transcriptomic and proteomic approaches. *Journal of Biomedicine and Biotechnology* 2011; 2011: 810242.
- [9] Munne S., Alikani M., Tomkin G., et al. Embryo morphology, developmental rates, maternal age are correlated with chromosome abnormalities. *Fertility and Sterility* 1995; 64: 382-91.
- [10] Fragouli E., Alfarawati S., Spath K., et al. The origin and impact of embryonic aneuploidy. *Hum. Genet.* 2013; 132: 1001-13.
- [11] Munne S., Chen S., Colls P., et al. Maternal age, morphology, development and chromosome abnormalities in over 6000 cleavage-stage embryos. *Reproductive Biomedicine Online* 2007; 14: 628-34.
- [12] Grondahl M.L., Christiansen S.L., Kesmodel U.S., et al. Effect of women's age on embryo morphology, cleavage rate and competence-A multicenter cohort study. *PLoS One* 2017; 12(4): E0172456.
- [13] Thornhill A., Geraedts J., Harper J.C., et al. ESHRE PGD Consortium «Best practice guidelines for PGD and PGS». *Human Reproduction* 2005; 20: 35-48.
- [14] Lalioti M. Can preimplantation genetic diagnosis overcome recurrent pregnancy failure *Current Opinion in Obstetrics and Gynecology* 2008; 20: 199-204.
- [15] Griffin D., Wilton L., Handyside A., et al. Dual fluorescent in situ hybridisation for simultaneous detection of X and Y chromosome-specific probes for the sexing of human preimplantation embryonic nuclei. *Human Genetics* 1992; 89: 18-22.
- [16] Munne S., Lee A., Rosenwaks Z., et al. Diagnosis of major chromosome aneuploidies in human preimplantation embryos. *Human Reproduction* 1993; 8: 2185-91.
- [17] Staessen C., Platteau P., Assche V., et al. Comparison of blastocyst transfer with or without preimplantation genetic diagnosis for aneuploidy screening in couples with advanced maternal age: A prospective randomized controlled trial. *Human Reproduction* 2004; 19: 2849-58.
- [18] Staessen C., Verpoest W., Donoso P., et al. Preimplantation genetic screening does not improve delivery rate in women under the age of 36 following single-embryo transfer. *Human Reproduction* 2008; 23: 2818-25.
- [19] Mastenbroek S., Twisk M., van Echten-Arends J., et al. In vitro fertilization with preimplantation genetic screening. *The New England Journal of Medicine* 2007; 357: 9-17.
- [20] Checa M., Alonso-Coello P., Sola I., et al. IVF/ICSI with or without preimplantation genetic screening for aneuploidy in couples without genetic disorders: A systematic review and meta-analysis. *Journal of Assisted Reproduction and Genetics* 2009; 26: 273-83.
- [21] Practice Committee of the Society for Assisted Reproductive technology and the Practice Committee of the American Society for Reproductive Medicine. Preimplantation genetic testing: A Practice Committee opinion. *Fertility and Sterility* 2008; 90: 136-43.
- [22] Harper J., Coonen E., De Rycke M., et al. What next for preimplantation genetic screening A position statement from the ESHRE PGD Consortium steering committee. *Human Reproduction* 2010; 25: 821-23.
- [23] Barbash-Hazan S., Frumkin T., Malcov M., et al. Preimplantation aneuploid embryos undergo self-correction in correlation with their developmental potential. *Fertility and Sterility* 2009; 92: 890-6.
- [24] Baart E.B., Martini E., Berg I., et al. Preimplantation genetic screening reveals a high incidence of aneuploidy and mosaicism in embryos from young women undergoing IVF. *Hum. Reprod.* 2006; 21(1): 223-33.
- [25] Los F.J., Opstal D., Berg C. The development of cytogenetically normal, abnormal and mosaic embryos: A theoretical model. *Hum. Reprod. Update* 2004; 10(1): 79-94.
- [26] Wilton L., Thornhill A., Traeger-Synodinos J., et al. The causes of misdiagnosis and adverse outcomes in PGD. *Hum. Reprod.* 2009; 24(5): 1221-8.
- [27] Wilton L. Preimplantation genetic diagnosis and chromosome analysis of blastomeres using comparative genomic hybridization. *Human Reproduction Update* 2005; 11: 33-41.
- [28] Kallioniemi A., Kallioniemi P., Sudar D., et al. Comparative genetic hybridisation for molecular cytogenetic analysis of solid tumors. *Science* 1992; 258: 818-21.
- [29] Wells D., Alfarawati S., Fragouli E. Use of comprehensive chromosomal screening for embryo assessment: Microarrays and CGH. *Molecular Human Reproduction* 2008; 12: 703-10.
- [30] Wells D., Sherlock J., Handyside A., et al. Detailed chromosomal and molecular genetic analysis of single cells by whole genome amplification and comparative genomic hybridisation. *Nucleic Acids Research* 1999; 27: 1214-8.
- [31] Voullaire L., Wilton L., Sargeant P., et al. Preimplantation aneuploidy screening using comparative genomic hybridisation or fluorescence in situ hybridisation of embryos from patients recurrent implantation failure. *Fertility and Sterility* 2003; 80: 860-8.

- [32] Shaffer L. Targeted genomic microarray analysis for identification of chromosome abnormalities in 1500 consecutive clinical cases. *The Journal of Pediatrics* 2006; 149(1): 98-102.
- [33] Shaffer L.G., Bui T.H. Molecular cytogenetic and rapid aneuploidy detection methods in prenatal diagnosis. *Am. J. Med. Genet. C Semin. Med. Genet.* 2007; 145: 87-98.
- [34] Handyside A.H. 24-chromosome copy number analysis: A comparison of available technologies. *Fertility and Sterility* 2013; 100(3): 595-602.
- [35] Treff N.R., Fedick A., Tao X., et al. Evaluation of targeted next-generation sequencing-based preimplantation genetic diagnosis of monogenic disease. *Fertil. Steril.* 2013; 99: 1377-84.
- [36] Fiorentino F., Bono S., Biricik A., et al. Application of next-generation sequencing technology for comprehensive aneuploidy screening of blastocysts in clinical preimplantation genetic screening cycles. *Human Reproduction* 2014; 29(12): 2802-13.
- [37] Zozola S., Schiewe M.C., Blazek J., et al. Reanalysis of failed vitrified euploid blastocyst transfer cycles using next-generation sequencing. *Fertility and Sterility* 2015; 104(3): 279.
- [38] Latham K., Garrels J., Chang C., et al. Analysis of embryonic mouse development: Construction of a high-resolution, two-dimensional gel protein database. *Applied and theoretical electrophoresis* 1992; 2: 163-70.
- [39] Navarrete S., Tonack S., Kirstein M., et al. Two insulin-responsive glucose transporter isoforms and the insulin receptor are developmentally expressed in rabbit preimplantation embryos. *Reproduction* 2004; 128: 503-16.
- [40] Wang Y., Puscheck E., Lewis J., et al. Increases in phosphorylation of SAPK/JNK and p38MAPK correlate negatively with mouse embryo development after culture in different media. *Fertility and Sterility* 2005; 1: 1144-54.
- [41] Gutstein H., Morris J., Annangudi S., et al. Microproteomics: Analysis of protein diversity in small samples. *Mass Spectrometry Reviews* 2008; 27: 316-30.
- [42] Katz-Jaffe M., Gardner D., Schoolcraft W. Proteomic analysis of individual human embryos to identify novel biomarkers of development and viability. *Fertility and Sterility* 2006; 85(1): 101-7.
- [43] Dominguez F., Gadea B., Esteban F., et al. Comparative protein-profile analysis of implanted versus non-implanted human blastocysts. *Human Reproduction* 2008; 23(9): 1993-2000.
- [44] Dominguez F., Meseguer M., Aparicio-Ruiz B., et al. New strategy for diagnosing embryo implantation potential by combining proteomics and timelapse technologies. *Fertility and Sterility* 2015; 104(4): 908-14.
- [45] O'Neill C. The role of PAF in embryo physiology. *Human Reproduction Update* 2005; 11: 215-28.
- [46] Cervero A., Horcajadas J., Dominguez F., et al. Leptin system in embryo development and implantation: A protein in search of a function. *Reproductive Biomedicine Online* 2005; 10: 217-23.
- [47] Mutz K.O., Heilkenbrinker A., Lonne M., et al. Transcriptome analysis using next-generation sequencing. *Current opinion in biotechnology* 2013; 24(1): 22-30.
- [48] Biase F., Everts R., Oliveira R., et al. Messenger RNAs in MII oocytes correlate with successful embryo development to the blastocyst stage. *Zygote* 2012; 10: 1-11.
- [49] Ouandaogo Z., Frydman N., Hesters L., et al. Differences in transcriptomic profiles of human cumulus cells isolated from oocytes at GV, MI and MII stages after in vivo and in vitro oocyte maturation. *Human Reproduction* 2012; 27: 2438-47.
- [50] McKenzie L., Pangas S., Carson S., et al. Human cumulus granulosa cell gene expression: A predictor of fertilization and embryo selection in women undergoing IVF. *Human Reproduction* 2004; 19: 2869-74.
- [51] Assou S., Haouzi D., De Vos J., et al. Human cumulus cells as biomarkers for embryo and pregnancy outcomes. *Molecular Human Reproduction* 2010; 16: 531-8.
- [52] Wood J., Dumesic D., Abbott D., et al. 3rd Molecular abnormalities in oocytes from women with polycystic ovary syndrome revealed by microarray analysis. *The Journal of Clinical Endocrinology and Metabolism* 2007; 92: 705-13.
- [53] Egea R.R., Puchalt N.G., Escrivá M.M., et al. OMICS: Current and future perspectives in reproductive medicine and technology. *J. Hum. Reprod. Sci.* 2014; 7(2): 73-92.
- [54] Zhang X., Jafari N., Barnes R., et al. Studies of gene expression in human cumulus cells indicate pentraxin 3 as a possible marker for oocyte quality. *Fertility and Sterility* 2005; 83 Suppl 1: 1169-79.
- [55] Inan M., Al-Hassan S., Ozand P., et al. Transcriptional profiling of granulosa cells from a patient with recurrent empty follicle syndrome. *Reproductive Biomedicine Online* 2006; 13: 481-91.
- [56] Assou S., Haouzi D., Mahmoud K., et al. A non-invasive test for assessing embryo potential by gene expression profiles of human cumulus cells: A proof of concept study. *Molecular Human Reproduction* 2008; 14: 711-9.
- [57] Brison D., Hollywood K., Arnesen R., et al. Predicting human embryo viability: The road to non-invasive analysis of the secretome using metabolic footprinting. *Reproductive Biomedicine Online* 2007; 15: 296-302.
- [58] Singh R., Sinclair K. Metabolomics: Approaches to assessing oocyte and embryo quality. *Theriogenology* 2007; 68 Suppl 1: 56-62.

- [59] Nagy Z., Sakkas D., Behr B. Symposium: Innovative techniques in human embryo viability assessment. Non-invasive assessment of embryo viability by metabolomic profiling of culture media. *Reproductive Biomedicine Online* 2008; 17: 502-7.
- [60] Borges E., Braga D., Setti A.S., et al. Non-invasive prediction of blastocyst implantation, ongoing pregnancy and live birth, by mass spectrometry lipid fingerprinting. *JBRA Assist. Reprod.* 2016; 20(4): 227-31.
- [61] Rubakhin S.S., Elena V., Romanova E.V., et al. Profiling Metabolites and Peptides in Single Cells. *Nat. Methods* 2011; 8 Suppl 4: 20-9.
- [62] Botros L., Sakkas D., Seli E. Metabolomics and its application for non-invasive embryo assessment in IVF. *Molecular Human Reproduction* 2008; 14(12): 679-90.
- [63] Renard J., Philippon A., Menezo Y. In vitro uptake of glucose by bovine blastocysts. *Journal of Reproduction and Fertility* 1980; 58(1): 161-4.
- [64] Van den Bergh M., Devreker F., Emilian S., et al. Glycolytic activity: A possible tool for human blastocyst selection. *Reproductive Biomedicine Online* 2001; 3 Suppl 1: 8.
- [65] Houghton F., Hawkhead J., Humpherson P., et al. Non-invasive amino acid turnover predicts human embryo developmental capacity. *Human Reproduction* 2002; 17(4): 999-1005.
- [66] Brison D., Houghton F., Falconer D., et al. Identification of viable embryos in IVF by non-invasive measurement of amino acid turnover. *Human Reproduction* 2004; 19(10): 2319-24.
- [67] Picton H., Elder K., Houghton F., et al. Association between amino acid turnover and chromosome aneuploidy during human preimplantation embryo development in vitro. *Molecular Human Reproduction* 2010; 16(8): 557-69.
- [68] Seli E., Sakkas D., Scott R., et al. Noninvasive metabolomic profiling of embryo culture media using Raman and near-infrared spectroscopy correlates with reproductive potential of embryos in women undergoing in vitro fertilization. *Fertility and Sterility* 2007; 88: 1350-7.
- [69] Li D., Zhai W., Li Y., et al. Recent progress in surface enhanced Raman spectroscopy for the detection of environmental pollutants. *Microchimica Acta* 2014; 181: 23-43.
- [70] Waddington C. The epigenotype. *International Journal of Epidemiology* 2012; 41: 10-3.
- [71] Singh K., Jaiswal D. Human male infertility: A complex multifactorial phenotype. *Reproductive Sciences* 2011; 18: 418-25.
- [72] Zama A., Uzumcu M. Epigenetic effects of endocrine-disrupting chemicals on female reproduction: An ovarian perspective. *Front. Neuroendocrinol.* 2010; 31: 420-39.
- [73] Iliadou A., Janson P., Cnattingius S. Epigenetics and assisted reproductive technology. *Journal of Internal Medicine* 2011; 270: 414-20.
- [74] van Montfoort A., Hanssen L., Sutter P., et al. Assisted reproduction treatment and epigenetic inheritance. *Human Reproduction Update* 2012; 18: 171-97.
- [75] Berger S. The complex language of chromatin regulation during transcription. *Nature* 2007; 447: 407-12.
- [76] Odom L., Segars J. Imprinting disorders and assisted reproductive technology. *Current Opinion in Endocrinology, Diabetes and Obesity* 2010; 17: 517-22.
- [77] Sato A., Otsu E., Negishi H., et al. Aberrant DNA methylation of imprinted loci in superovulated oocytes. *Human Reproduction* 2007; 22(1): 26-35.
- [78] Montfoort A.P., Hanssen L.L., de Sutter P., et al. Assisted reproduction treatment and epigenetic inheritance. *Hum. Reprod. Update* 2012; 18(2): 171-97.
- [79] Bunkar N., Pathak N., Lohiya N.K., et al. Epigenetics: A key paradigm in reproductive health. *Clin. Exp. Reprod. Med.* 2016; 43(2): 59-81.
- [80] Huang J.C., Lei Z.L., Shi L.H., et al. Comparison of histone modifications in in vivo and in vitro fertilization mouse embryos. *Biochem. Biophys. Res. Commun.* 2007; 354(1): 77-83.
- [81] Ma H., Marti-Gutierrez N., Park S.W., et al. Correction of a pathogenic gene mutation in human embryos. *Nature* 2017; 548: 413-9.
- [82] Liang P., Xu Y., Zhang X., et al. CRISPR/Cas9-mediated gene editing in human tripronuclear zygotes. *Protein Cell* 2015; 6(5): 363-72.
- [83] Ledford H. Where in the world could the first CRISPR baby be born *Nature* 2015; 526: 310-1.